

Randomly dilute Ising model: A nonperturbative approach

Matthieu Tissier¹, Dominique Mouhanna¹, Julien Vidal², and Bertrand Delamotte¹

¹ *Laboratoire de Physique Théorique et Hautes Energies, CNRS UMR 7589,*

Université Pierre et Marie Curie Paris 6, 4 place Jussieu, 75252 Paris Cedex 05 France

² *Groupe de Physique des Solides, CNRS UMR 7588,*

Universités Pierre et Marie Curie Paris 6 et Denis Diderot Paris 7, 2 place Jussieu, 75251 Paris Cedex 05 France

The N -vector cubic model relevant, among others, to the physics of the randomly dilute Ising model is analyzed in arbitrary dimension by means of an exact renormalization-group equation. This study provides a unified picture of its critical physics between two and four dimensions. We give the critical exponents for the three-dimensional randomly dilute Ising model which are in good agreement with experimental and numerical data. The relevance of the cubic anisotropy in the $O(N)$ model is also treated.

PACS numbers: 75.10.Nr, 11.10.Hi, 11.15.Tk, 64.60.-i

Understanding the effects of a weak quenched disorder on magnetic systems is one of the great challenges of statistical physics. In this context, much attention has been paid to the Ising model with randomly distributed non-magnetic impurities, i.e. the random Ising (RI) model (see¹ and² for reviews) described by the following Hamiltonian :

$$H = -J \sum_{\langle i,j \rangle} \varepsilon_i \varepsilon_j S_i S_j \quad (1)$$

with $J > 0$. In Eq. (1), the S_i are Ising variables and the ε_i are quenched, uncorrelated, random variables taking the value 1 with probability p and 0 with probability $1 - p$. According to the Harris criterion³, the critical behavior of a pure system is altered by the disorder if its specific-heat exponent α is positive, as it is the case for the three-dimensional Ising model for which $\alpha \simeq 0.109$, $\beta \simeq 0.326$, $\gamma \simeq 1.24$, $\nu \simeq 0.630$ and $\eta \simeq 0.0335$ (see⁴).

Experimentally, there now exist several compounds corresponding to a RI model. They essentially consist in crystalline difluoride of iron or manganese where FeF_2 or MnF_2 is replaced by a nonmagnetic material such as ZnF_2 . Experiments performed on $\text{Fe}_p\text{Zn}_{1-p}\text{F}_2$ or $\text{Mn}_p\text{Zn}_{1-p}\text{F}_2$ with different values of the concentration p have revealed scaling laws with exponents clearly distinct from those of the pure Ising model. One has (see²), for instance, for $\text{Fe}_p\text{Zn}_{1-p}\text{F}_2$: $-0.12 \leq \alpha \leq -0.06$, $0.34 \leq \beta \leq 0.38$, $1.28 \leq \gamma \leq 1.4$ and $0.68 \leq \nu \leq 0.72$.

Numerically, early Monte Carlo simulations had confirmed the relevance of disorder for the Ising model (see²). The situation was however unclear since the critical exponents seemed to vary with p . Recently, Ballesteros et al.⁵, by a careful infinite volume extrapolation of the data based upon an analysis of the correction-to-scaling terms, have reached the conclusion of dilution-independent critical exponents : $\alpha = -0.051(16)$, $\beta = 0.3546(28)$, $\gamma = 1.342(10)$, $\nu = 0.6837(53)$ and $\eta = 0.0374(45)$ for $0.4 \leq p \leq 0.9$.

From a theoretical point of view, the continuous field theory relevant to the study of the RI model is obtained by considering N replica of the model (1) and by taking

the average over all possible realizations of the Gaussian distributed disorder. This leads to the Ginzburg-Landau-Wilson action for the N -vector model with cubic anisotropy^{6,7} :

$$S = \int d^d x \left\{ \frac{1}{2} \sum_{a=1}^N [(\partial_\mu \phi_a)^2 + m^2 \phi_a^2] + u_1 \left(\sum_{a=1}^N \phi_a^2 \right)^2 + u_2 \sum_{a=1}^N \phi_a^4 \right\} \quad (2)$$

with $u_1 \propto -(1-p) < 0$ and $u_2 > 0$. The physics of the RI model is then obtained through the $N \rightarrow 0$ limit of the renormalization group (RG) equations obtained from action (2). The determination of these RG equations has been the subject of an intense activity, mainly performed within perturbative frameworks. Different renormalization schemes have been used : minimal subtraction (MS) scheme^{6,8,9,10,11} or massive scheme^{12,13,14,15,16,17} directly in three dimensions where the β functions have been computed up to five¹¹ and six loops¹⁵ respectively. In each scheme, two different kinds of expansions for the corresponding β functions have been considered : *i*) ϵ -expansion (actually $\sqrt{\epsilon}$ -expansion)^{6,8,9,10,11} in the MS scheme and pseudo- $\sqrt{\epsilon}$ -expansion¹⁸ in the massive scheme *ii*) expansion in the coupling constants in three dimensions^{12,13,14,15,16,17}. It has, however, appeared that the series obtained in case *i*) are non-Borel summable and thus inappropriate for the quantitative evaluation of critical exponents behavior^{9,10,19}, a result anticipated by several authors^{20,21} within an analysis of the zero-dimensional problem (see however²²). In case *ii*), generalizations of Padé-Borel resummation techniques for several variables have been considered. In the MS scheme, Chisholm-Borel resummation scheme has been used^{9,19,23}. This method has led to satisfying results at low order but has also appeared to be unstable with respect to the order of approximation. Indeed, the five-loop calculation does not allow to obtain real coordinates for the RI fixed point. This point has been related to the possible Borel-non-summability of the series^{2,19,23}. In

the massive scheme, Pelissetto and Vicari¹², using a procedure of resummation formulated within a subsequent analysis of the zero-dimensional problem²², have employed Padé-Borel and conformal Padé-Borel resummation techniques within a six-loop RG calculation. They found a fixed point with exponents $\gamma = 1.330(17)$, $\nu = 0.678(10)$, $\eta = 0.030(3)$ that are in rather good agreement with previous perturbative results^{13,14,16} and with the present-day numerical data. Let us however mention that, even in this case, the location of the fixed point is found to be calculation-dependent. This has led Pelissetto et al. to consider the possibility of an optimal truncation, possibly shorter than that considered in¹², a fact that could constitute a limitation of the method.

Despite the good agreement between experimental and numerical data and the perturbative results, we would like to stress upon the need for a complementary approach, not relying on perturbative expansions and this, for several reasons. The first one is that the perturbative field theoretical treatment implies the use of sophisticated resummation techniques which, if they proved to lead to safe results in the case of pure magnets, are far from being under control in the case of disordered systems^{2,20,21,22}. The second reason for the use of a non-perturbative treatment is that there exist cases where the perturbative approach apparently seems to provide incorrect results. For instance, in the context of frustrated magnets, six-loop perturbative RG computations predict the existence of a fixed point²⁴ and, thus, a universal behavior for their critical physics. The experimental data, although displaying scaling laws, are rather in favor of a nonuniversal behavior, the critical exponents varying from one material to the other. Computations based on the exact RG formalism have shown to be able to reproduce this phenomenology^{25,26}. Finally, there are situations where only specific techniques work well. For example, in two dimensions, one can use integrability or conformal invariance which, however, only operate in this dimension. All these features plead toward the use of a method that does not rely on the treatment of divergent series, provides a systematic way to improve the results and is able to tackle with the physics in any dimension.

A Wilson-like²⁷ nonperturbative treatment of (2) has been first performed by Newman and Riedel²⁸ on the basis on the scaling-field (SF) method. However, due to the complexity of its practical implementation, this approach has been given up and perturbative approaches largely preferred. We used here a nonperturbative approach based on the concept of effective average action $\Gamma_k[\phi]$ ^{29,30} which appears as a very efficient tool. The quantity $\Gamma_k[\phi]$ has the meaning of a coarse-grained free energy at the scale k in the sense that it only includes the fluctuations with momenta $q \geq k$. It is a function of the effective order parameter ϕ at the scale k , the analog of a magnetization at this scale. In the low-energy limit $k \rightarrow 0$, all modes are integrated out so that $\Gamma_{k=0}$

identifies with the usual effective action Γ . At the microscopical scale Λ , no fluctuation has been integrated and $\Gamma_{k=\Lambda}$ identifies with the microscopic action S . The k -dependence of Γ_k is controlled by an exact evolution equation³¹:

$$\frac{\partial \Gamma_k}{\partial t} = \frac{1}{2} \text{Tr} \left\{ (\Gamma_k^{(2)} + R_k)^{-1} \frac{\partial R_k}{\partial t} \right\}, \quad (3)$$

where $t = \ln(k/\Lambda)$. The trace has to be understood as a momentum integral as well as a summation over internal indices. In Eq.(3), $\Gamma_k^{(2)}$, the second derivative of Γ_k , is the *exact field-dependent* inverse propagator and R_k is the running infrared cutoff that suppresses the propagation of modes with momenta $q < k$. A convenient cutoff is provided by³⁰: $R_k(q) = Zq^2/[\exp(q^2/k^2)-1]$, where Z is the field renormalization. The effective average action Γ_k is a functional invariant under the symmetry group of the system and thus includes all powers of all invariants – and their derivatives – built out from the average order parameter. Thus, Eq.(3) is a nonlinear functional equation, too difficult to be solved exactly so that, in practice, one has to deal with a truncated form of Γ_k . We choose here to perform an expansion of Γ_k around its minimum, keeping a finite number of monomials in the invariants and including the most relevant derivative terms³⁰. We first consider the following truncation:

$$\Gamma_k = \int d^d x \left\{ \sum_{a=1}^N \frac{Z}{2} (\partial_\mu \phi_a)^2 + u_1 \left[\sum_{a=1}^N (\phi_a^2 - \kappa) \right]^2 + u_2 \sum_{a=1}^N (\phi_a^2 - \kappa)^2 \right\} \quad (4)$$

where $\{u_1, u_2, \kappa, Z\}$ are the k -dependent coupling constants that parametrize the evolution of the model with the scale k . The action (4) displays a discrete $\mathbb{Z}_N$ permutation symmetry. For $Nu_1 + u_2 > 0$, the minimum of action (4) is given by a replica symmetric configuration defined by $\phi_a^{min} = \sqrt{\kappa}$ for all a . Expression (4) thus corresponds to a quartic expansion of Γ_k around the minimum ϕ_a^{min} . There is no residual symmetry in the minimum so that the symmetry breaking scheme is given by a fully broken $\mathbb{Z}_N$ group. The spectrum of the theory in the broken phase is given by the eigenvalues of $\Gamma_k^{(2)}$ at the minimum and consists in a singlet of mass $m_1^2 = 8\kappa(Nu_1 + u_2)$ and a $(N-1)$ uplet of mass $m_2^2 = 8\kappa u_2$. When $u_2 = 0$, this $(N-1)$ uplet corresponds to the Goldstone modes of the $O(N)$ -invariant model that gets here a mass proportionnal to the $O(N)$ -symmetry-breaking-coupling-constant u_2 .

Using Eq.(3) and Eq.(4), we have derived the flow equations for the dimensionless renormalized coupling constants κ_r , u_{1r} , u_{2r} , and Z :

$$\begin{aligned}
\partial_t \kappa_r &= -(d-2+\eta)\kappa_r - \frac{2v_d}{N(Nu_{1r}+u_{2r})} \left[3(Nu_{1r}+u_{2r})L_1^d[m_{1r}^2] + (N-1)(Nu_{1r}+3u_{2r})L_1^d[m_{2r}^2] \right] \\
\partial_t u_{1r} &= (d-4+2\eta)u_{1r} - \frac{2v_d}{N^2\kappa_r} \left[-Nu_{1r}L_1^d[m_{1r}^2] + Nu_{1r}L_1^d[m_{2r}^2] + 36\kappa_r(Nu_{1r}+u_{2r})^2L_2^d[m_{1r}^2] \right. \\
&\quad \left. + \left(4(N-1)N^2\kappa_ru_{1r}^2 + 24N(N-1)\kappa_ru_{1r}u_{2r} + 36\kappa_ru_{2r}^2 \right) L_2^d[m_{2r}^2] - 8\kappa_r(Nu_{1r}+3u_{2r})^2L_{1,1}^d[m_{1r}^2, m_{2r}^2] \right] \\
\partial_t u_{2r} &= (d-4+2\eta)u_{2r} - \frac{2v_d}{N\kappa_r} \left[Nu_{1r}L_1^d[m_{1r}^2] - Nu_{1r}L_1^d[m_{2r}^2] + 36(N-2)\kappa_ru_{2r}^2L_2^d[m_{2r}^2] \right. \\
&\quad \left. + 8\kappa_r(Nu_{1r}+3u_{2r})^2L_{1,1}^d[m_{1r}^2, m_{2r}^2] \right] \\
\eta &= -\partial_t \ln Z = -\frac{256\kappa_r v_d}{Nd} \left[(N-1)u_{2r}^2M_{2,2}^d[0, m_{2r}^2] + (Nu_{1r}+u_{2r})^2M_{2,2}^d[0, m_{1r}^2] \right]
\end{aligned} \tag{5}$$

where $1/v_d = 2^{d+1}\pi^{d/2}\Gamma(d/2)$. In Eqs.(5), $L_{n_1, n_2}^d[m_{1r}^2, m_{2r}^2]$, $L_n^d[m_{ir}^2]$ and $M_{n_1, n_2}^d[m_{1r}^2, m_{2r}^2]$ are the “threshold functions”³² that encode the nonperturbative aspects of the RG equations (5). We show how this set of equations provides a qualitative and, partly, quantitative description of the physics of model (2) between $d=2$ and $d=4$.

Around $d=4$, the fixed point coupling constants u_{1r} and u_{2r} are expected to be of order $\epsilon = 4-d$. One can thus obtain simplifications of Eqs.(5) performing a small mass $-m_{1r}$ and m_{2r} - expansion of the threshold functions. Using their asymptotic behavior one finds $\eta = O(u_{1r}^2, u_{2r}^2)$ and :

$$\partial_t u_{1r} = -\epsilon u_{1r} + \frac{1}{2\pi^2} [(N+8)u_{1r}^2 + 6u_{1r}u_{2r}] \tag{6}$$

$$\partial_t u_{2r} = -\epsilon u_{2r} + \frac{1}{2\pi^2} [9u_{2r}^2 + 12u_{1r}u_{2r}] . \tag{7}$$

These equations are those obtained directly in the weak-coupling expansion of the N -vector cubic model^{6,7}. For $N \neq 0$ they display three nontrivial fixed points : the $O(N)$ fixed point ($u_{1r}^* > 0, u_{2r}^* = 0$), the N -decoupled Ising one ($u_{1r}^* = 0, u_{2r}^* > 0$) and the cubic one ($u_{1r}^* > 0, u_{2r}^* \neq 0$). The cubic fixed point is stable for $N > N_c$ and unstable for $N < N_c$ with $N_c(d=4-\epsilon) = 4-2\epsilon + O(\epsilon^2)$, as known from early perturbative approaches^{6,7}. For $N=0$ the only fixed point of the equations (6) and (7) is the Gaussian one but this is an artefact of the leading order of perturbation theory that disappears at second order in ϵ . This leads to the famous $\sqrt{\epsilon}$ -expansion^{6,8}. However, when the Eqs.(5), obtained with the simplest ansatz (4), are analyzed without ϵ -expansion, the random fixed point already appears. All the fixed points found here persist when the dimension is lowered from $d=4$ to the vicinity of $d=2$, a case that deserves a special treatment³³.

Around $d=2$, one also expects the Eqs.(5) to well describe, even quantitatively, the physics in the vicinity of the $O(N > 1)$ fixed-point. The reason is that, around

this dimension, the phase transition associated to this fixed point takes place at a temperature of order $\epsilon = d-2$. Since the anomalous dimension is of the same order of magnitude, this situation must be well described even with our wild truncation that neglects high order derivative terms. Around $d=2$, $1/(N\kappa_r) = T_r$ provides the effective perturbative temperature of the associated $O(N)$ non-linear sigma (NL σ) model. One also has to consider u_2 as a small perturbation around the $O(N)$ fixed point. Thus, the procedure now consists in a double expansion of Eqs.(5) in the large mass m_{1r} and in the small coupling constant u_{2r} . Using the asymptotic behaviors of the threshold functions, one obtains : $\eta = T_r/(2\pi N)$ and :

$$\partial_t T_r = \epsilon T_r - \frac{(N-2)}{2\pi} T_r^2 \tag{8}$$

$$\partial_t u_{1r} = -2u_{1r} + 4u_{1r}^2 \frac{(N-1)}{\pi} \ln 2 \tag{9}$$

$$\partial_t u_{2r} = \left(-2 + \epsilon + \frac{4}{N\pi} T_r \right) u_{2r} . \tag{10}$$

These equations admit a nontrivial fixed point with $T_r^* = 2\pi\epsilon/(N-2)$, $u_{1r}^* = \pi/(2(N-1)\ln 2)$ and $u_{2r}^* = 0$ which corresponds to the standard $O(N)$ fixed point of the NL σ model. It is then interesting to consider the flow of u_{2r} around this fixed point. One has :

$$\partial_t \delta u_{2r} = \left(-2 + \frac{(N+6)}{(N-2)} \epsilon \right) \delta u_{2r} . \tag{11}$$

This shows that the cubic perturbation is relevant for $N > N_c$ and irrelevant for $N < N_c$ with $N_c(d=2+\epsilon) = 2+4\epsilon + O(\epsilon^2)$, in agreement with the perturbative approach of the NL σ model around $d=2$ ³⁴. Concerning the fixed points with discrete symmetry (Ising $O(1)$, N -decoupled Ising, cubic and random Ising), they cannot, around $d=2$, be properly described by our truncation. Actually, the large values of η at these fixed points around $d=2$ likely require the inclusion in (4) of all powers of the field and high order derivative terms.

In $d = 3$, in order to obtain satisfying quantitative results, we have pushed the calculation using a more sophisticated ansatz than (4). We have considered all monomials up to order eight in the fields as well as all derivative terms including two derivatives and four fields and established the RG equations for the corresponding coupling constants. These equations are too long to be given here³⁵ and we just give our results. First, we have tested our calculation on the $O(N)$ model. In the Ising and Heisenberg cases, we have found $\nu = 0.627$ and $\nu = 0.719$, respectively, which are very reasonable compared to previous results (see⁴). The values of the anomalous dimensions $\eta = 0.053$ and $\eta = 0.0452$ are less convincing and related to the omission of higher order derivative terms in Γ_k . Considering the cubic anisotropy, we have determined an estimate of N_c above which the cubic fixed point is stable and found $N_c(d = 3) = 3.1$. This value differs significantly from those obtained from recent perturbative approaches for which $N_c < 3$ ^{10,11,14,15,16,17,18}. For instance, in a recent six-loop calculation, N_c is found to be equal to $2.89(4)$ ¹⁵, a value already obtained from lower order computations^{16,17}. If this last result is con-

firmed, our result should be imputed to the poor determination of η . The strong dependence of N_c with respect to η already appears in a hierarchical approximation where, neglecting any field renormalization, $N_c = 2.219$ ³⁶. Note also that the result $N_c = 3.4$ was found within the work of Newman and Riedel²⁸. For the RI fixed point we have found $\nu = 0.67$ and $\eta = 0.05$ that give, through scaling laws, $\gamma = 1.306$, $\beta = 0.351$ and $\alpha = -0.01$. These results are in good agreement with perturbative methods^{2,9,12,13,16,17,19?} and also with numerical estimates, while they show a certain discrepancy with the SF method that provides $\nu = 0.697$, $\gamma = 1.39$ and $\eta = 0.015$ ²⁸.

Finally, we indicate that we have noted a rapid convergence of our results with respect to the order of the truncation in the fields. However, a challenging and still open question is that of the convergence when including higher order derivative terms which is essential for an improvement of the anomalous dimension.

We thank M. L. Rosinberg, B. N. Shalaev and G. Tarjus for discussions.

-
- ¹ R. B. Stinchcombe, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz, (Academic Press, New York, 1983), Vol. 7, p.152 .
 - ² R. Folk and Yu. Holovatch and T. Yavors'kii, cond-mat/0106468 .
 - ³ A. B. Harris, J. Phys. C **7**, 1671 (1974).
 - ⁴ A. Pelissetto and E. Vicari, cond-mat/0012164 .
 - ⁵ H. G. Ballesteros, L. A. Fernandez, V. Martin-Mayor, A. Munoz Sudupe, G. Parisi, and J. J. Ruiz-Lorenzo, Phys. Rev. B **58**, 2740 (1998).
 - ⁶ G. Grinstein and A. Luther, Phys. Rev. B **13**, 1329 (1976).
 - ⁷ A. Aharony, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz, (Academic Press, New York, 1976), Vol. 6, p.357 .
 - ⁸ D. E. Khmel'nitskii, Sov. Phys. JETP **41**, 981 (1975).
 - ⁹ R. Folk, Yu. Holovatch, and T. Yavors'kii, JETP Lett. **69**, 747 (1999).
 - ¹⁰ B. N. Shalaev, S. A. Antonenko, and A. I. Sokolov, Phys. Lett. A **230**, 105 (1997).
 - ¹¹ H. Kleinert and V. Schulte-Frohlinde, Phys. Lett. B **342**, 284 (1995).
 - ¹² A. Pelissetto and E. Vicari, Phys. Rev. B **62**, 6393 (2000).
 - ¹³ D. V. Pakhnin and A. I. Sokolov, Phys. Rev. B **61**, 15130 (2000).
 - ¹⁴ I. O. Mayer, A. I. Sokolov, and B. N. Shalaev, Ferroelectrics **95**, 93 (1989).
 - ¹⁵ J. M. Carmona, A. Pelissetto, and E. Vicari, Phys. Rev. B **61**, 15136 (2000).
 - ¹⁶ K. B. Varnashev, Phys. Rev. B **61**, 14660 (2000).
 - ¹⁷ K. B. Varnashev, J. Phys. A **33**, 3121 (2000).
 - ¹⁸ R. Folk, Yu. Holovatch, and T. Yavors'kii, Phys. Rev. B **62**, 12195 (2000).
 - ¹⁹ R. Folk, Yu. Holovatch, and T. Yavors'kii, Phys. Rev. B **61**, 15114 (2000).
 - ²⁰ A. J. McKane, Phys. Rev. B **49**, 12003 (1994).
 - ²¹ A. J. Bray, T. McCarthy, M. A. Moore, J. D. Reger, and A. P. Young, Phys. Rev. B **36**, 2212 (1987).
 - ²² G. Alvarez, V. Martin-Mayor, and J. J. Ruiz-Lorenzo, J. Phys. A **33**, 841 (2000).
 - ²³ R. Folk, Yu. Holovatch, and T. Yavors'kii, J. Phys. Stud. **2**, 213 (1998).
 - ²⁴ A. Pelissetto, P. Rossi, and E. Vicari, Phys. Rev. B **63**, 140414 (2001).
 - ²⁵ M. Tissier, B. Delamotte, and D. Mouhanna, Phys. Rev. Lett. **84**, 5208 (2000).
 - ²⁶ M. Tissier, B. Delamotte, and D. Mouhanna, cond-mat/0107183 .
 - ²⁷ K. G. Wilson and J. Kogut, Phys. Rep. C **12**, 75 (1974).
 - ²⁸ K. E. Newman and E. K. Riedel, Phys. Rev. B **25**, 264 (1982).
 - ²⁹ C. Wetterich, Nucl. Phys. B **352**, 529 (1991).
 - ³⁰ J. Berges, N. Tetradis, and C. Wetterich, hep-ph/0005122 .
 - ³¹ C. Wetterich, Phys. Lett. B **301**, 90 (1993).
 - ³² N. Tetradis and C. Wetterich, Nucl. Phys. B : Field Theory Stat. Syst. [FS] **422**, 541 (1994).
 - ³³ B. N. Shalaev, Phys. Rep. **237**, 129 (1994).
 - ³⁴ R. A. Pelcovits and D. R. Nelson, Phys. Lett. A **57**, 23 (1976).
 - ³⁵ M. Tissier, D. Mouhanna, J. Vidal, and B. Delamotte, in preparation .
 - ³⁶ K. Pinn, M. Rehwald, and C. Wiecekowsky, J. Stat. Phys. **95**, 1 (1999).